



## **DRY GRANULAR FERTILISER ENZYME ADDITIVE**

### **ACTIVE INGREDIENTS:**

**Phosphatase >250 mU/g**  
**Mannanase >1.25 kU/g**



**WARNING:**  
May Cause Allergic Skin Reaction

Also Contains

<0.01% reaction mass of 5-chloro-2-methyl-4-iso-thiazolin-3- one  
and 2-methyl-2H-isothiazol-3- one  
<0.1% 1,2-Benzisothiazol-3(2H)-one

Dry enzyme additive that enhances availability of soil-bound nutrients, in particular phosphorus, to plant roots and native soil microbes resulting in improved soil and plant health, which promotes plant growth and yield.



**50 GRAMS**

**Storage & Disposal:**

Store in original sachet in a cool, well-ventilated area, out of direct sunlight, avoiding temperature extremes.

**DO NOT** reuse this sachet for any other purpose.

Deliver to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation, and tree roots. Do not burn empty sachets or product.

**For further information refer to the Safety Data Sheet (SDS).**

Magno® is a registered trademark of Elemental Enzymes Australia Pty Ltd.

**Manufactured by:**

Elemental Enzymes Head Office,  
1685 Galt Industrial Blvd, St Louis,  
MO, 63132, USA.

t: +1 314 429 2885

e: [info@elementalenzymes.com](mailto:info@elementalenzymes.com)

w: [www.elementalenzymes.com](http://www.elementalenzymes.com)

**Distributed by:**

Elemental Enzymes Australia Pty Ltd  
40-42 Scott Street

Dandenong Victoria 3175

t: 1800 691 440

e: [info@elementalenzymes.com](mailto:info@elementalenzymes.com)

w: [www.elementalenzymes.com.au](http://www.elementalenzymes.com.au)

EE Magno GENERIC Leaflet  
ID:  
66x90mm

## DIRECTIONS FOR USE:

Magno® is for soil application only.

Magno® is designed for use on all soil types.

Magno® can be applied to dry soil if planting dry and will be available when the seed germinates.

### Mixing & Compatibility:

Check chemical mixture compatibility using a jar test using all products in proper proportion to establish physical compatibility prior to application.

This product is non-phytotoxic and compatible with many commonly used in furrow treatments including fertilisers, insecticides, pesticides, fungicides, micronutrients, and bio-stimulants. Can be mixed with most NPK fertilisers with a pH between 5 and 9, well as water carrier solution. Half fill mix tank with water and begin agitation. Add products in this order, mixing thoroughly after each addition: pesticides, fertilisers, Magno and finally adjuvants. Fill tank with remainder of water and continue agitation until solution is completely mixed.

SITUATION	USE PATTERN	RATE	CRITICAL COMMENTS
Horticultural crops	At planting, in-furrow with starter fertiliser or water (or both) for all horticultural crops.	15 to 50 g/ha	Apply as a total in-furrow spray volume of 30L to 140 L/ha using fertiliser and/or water carrier. Refer to mixing and compatibility statements for order of mixing.
	Vegetative stages through dripline application with fertiliser or water (or both) for all horticultural crops.	15 to 50 g/ha	Dilute in sufficient application volume to apply through drip or micro-sprinkler irrigation. Apply a sufficient quantity of irrigation to extend product through the root zone. Refer to mixing and compatibility statements for order of mixing.
	For side-dressing when applied within 15 cm of the plants.	15 to 50 g/ha	When side-dressing apply with liquid fertiliser solution in spray volume of 30L to 140 L/ha. Refer to mixing and compatibility statements for order of mixing.
Sugarcane	Both plant and ratoon.	10 to 20 g/ha	Apply at planting or on ratoon cane when stool splitting occurs to apply nutrition. Apply in a spray volume of 50 – 300 L/ha.

Prior to the addition of Magno® to the spray tank it is preferable to pre-dissolve using the following instructions. Fill a bottle or jug with approximately 1 litre of water for every hectare to be treated. Add pre-weighed Magno® at the suggested use rate. Close the container and shake thoroughly for 15 seconds; allow the solution to rest for 1 minute and then mix again. Repeat mixing and resting until Magno is fully dissolved.

### Product Use:

Apply Magno® in the root zone of target crops where it will act by release of both sugars from mucilage secreted by root tips as well as plant available orthophosphate from soil borne organic phosphates. The combined activity enhances nutrient uptake by the plant and increases microbial activity around the roots.



**Storage & Disposal:** Store in original sachet in a cool, well-ventilated area, out of direct sunlight, avoiding temperature extremes. DO NOT reuse this sachet for any other purpose. Deliver to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation, and tree roots. Do not burn empty sachets or product. **Safety Directions:** Avoid breathing dust. In case of inadequate ventilation wear respiratory protection. Wear protective gloves. Contaminated work clothing should not be allowed out of the workplace and must be washed before reusing. Wash hands thoroughly after handling. **First Aid:** IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs: Get medical advice/attention. **For further information refer to the Safety Data Sheet (SDS).**

**Manufactured by:**

Elemental Enzymes Head Office,  
1685 Galt Industrial Blvd, St Louis, MO,  
63132, USA.  
t: +1 314 429 2885  
e: [info@elementalenzymes.com](mailto:info@elementalenzymes.com)  
w: [www.elementalenzymes.com](http://www.elementalenzymes.com)

**Distributed by:**

Elemental Enzymes Australia Pty Ltd  
40-42 Scott Street  
Dandenong Victoria 3175  
t: 1800 691 440  
e: [info@elementalenzymes.com](mailto:info@elementalenzymes.com)  
w: [www.elementalenzymes.com.au](http://www.elementalenzymes.com.au)

Magno® is a registered trademark of Elemental Enzymes Australia Pty Ltd.  
Patent No. AU 2017233034

**Batch Number:** see packaging

**Date of Manufacture:** see packaging

**Expiry date:** 2 years from date of manufacture